

Ibuprofen (Motrin, Advil)

****DO NOT GIVE TO AN INFANT UNDER AGE OF 6 MONTHS****

Consult with a physician or your local pharmacist before giving with any other medications. Some medications may contain additional pain/fever reducing agents. Aspirin is not to be given to a child.

****Ibuprofen can be given every 6-8 hours****

DO NOT GIVE UNDER 6 MONTHS OF AGE	6-11 LBS	12-17 LBS	18-23 LBS	24-35 LBS	36-47 LBS	48-59 LBS	60-71 LBS	72-95 LBS
SUSPENSION 50MG/125ML	X	1.25 mL	1.875 mL	2.5 mL	3.75 mL	X	X	X
SUSPENSION 100MG/5ML	X	X	X	5 mL	7.5 mL	10 mL	12.5 mL	15 mL
CHEW-ABLE TABLETS (100 MG)	X	X	X	1 Tablet	1 ½ Tablets	2 Tablets	2 ½ Tablets	3 Tablets
Tablets (200 MG)	X	X	X	X	X	1 Tablet	1 ½ Tablets	2 ½ Tablets

Acetaminophen (Tylenol) Dose

*[12-22-2011] The U.S. Food and Drug Administration (FDA) changed policy and informed the public that an additional concentration of liquid acetaminophen marketed for "infants" (160 mg/5 mL) is now available at local stores. Only other liquid acetaminophen marketed for infants was only available in 80 mg/0.8 mL or 80 mg/mL concentrations. This change in the concentration could affect the amount of liquid given to an infant, and should be especially noted if someone is accustomed to using the 80 mg /0.8 mL or 80 mg/mL concentrations of liquid acetaminophen. *

**** All store shelves should now have the (160mg/5mL) Dosing****

**** PLEASE only use the dropper or dosage cup that comes with your medication to avoid overdosing. Once Bottle is complete throw bottle, dropper or dosage cup away to prevent over dosing of other medications****

Please follow the table below for dosage and do not give a higher dose unless directed by a health care provider ~ Aspirin should not be given to your child, **DO NOT GIVE ACETAMINOPHEN (TYLENOL) TO INFANTS UNDER 2 MONTHS OF AGE**, if less than 2 months of age and has a fever, seek medical attention and/or treatment.

Acetaminophen can be given every 4-6 hours

DO NOT GIVE UNDER 2 MONTHS OF AGE	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
160mg/5mL	1.25 mL	2.5 mL	3.75 mL	5 mL	7.5 mL	10 mL	12.5 mL	15 mL
Chew-able jr 160 mg tablets	X	X	X	1 Tablet	1 ½ Tablets	2 Tablets	2 ½ Tablets	3 Tablets
Suppository 80mg (6m-36m)	X	1 Supp.	1 ½ Supp.	2 Supp.	X	X	X	X
Suppository 120 mg (3-6 years)	X	X	1 Supp.	1 ½ Supp.	X	X	X	X
Suppository 325mg (6-12 Years)	X	X	X	½ Supp.	¾ Supp.	1 Supp.	X	2 Supp.